

Metabolism and Pharmacokinetics of Alternate Drinking Water Disinfectants

by Mohamed S. Abdel-Rahman,* Daniel Couri† and Richard J. Bull††

The chlorination of surface waters is known to elevate trihalomethanes; consequently, chlorine dioxide (ClO_2) is being considered as an alternative disinfectant. The primary products resulting from ClO_2 disinfection of waters are chlorites (ClO_2^-) and chlorates (ClO_3^-). Studies in rats revealed that ClO_2 is converted to chloride (Cl^-), ClO_2^- and ClO_3^- . ClO_2^- and ClO_3^- are excreted as Cl^- , ClO_2^- and Cl^- , ClO_2^- , ClO_3^- , respectively. Radioactivity was rapidly absorbed from the gastrointestinal tract following the administration of $^{36}\text{ClO}_2$ orally, and the half-life for the elimination of ^{36}Cl from the rat was 44 hr, corresponding to a rate constant of 0.016/hr. After 72 hr, radioactivity was highest in plasma, followed by kidney, lung, and stomach. ^{36}Cl in plasma reached a peak at 2 hr and 1 hr after oral administration of $^{36}\text{ClO}_2^-$ and $^{36}\text{ClO}_3^-$, respectively. ^{36}Cl excretion was greatest 24 hr after the administration of $^{36}\text{ClO}_3^-$, but in the case of $^{36}\text{ClO}_2^-$, the excretion probably represented saturation of the biotransformation and excretion pathway. A low activity in packed cells compared to plasma was detected in chlorate ingestion, rather than an even distribution in chlorite treatment. Chloroform determinations in rat blood after one year indicated that chloroform was significantly higher than control in the 100 and 1000 mg/l. ClO_2 groups. However, no significant values were observed in the 1 or 10 mg/l. ClO_2 and ClO_2 metabolites group. ClO_2 and its metabolites are eliminated from the body more rapidly than chlorine, and they do not appear to increase trihalomethane concentrations at low dosages.

Introduction

Chlorination is, and always has been, the most widely used method for disinfection of public water supplies. Concern over its possible adverse effects and those of its by-products on humans has increased considerably in recent years. The formation of trihalomethanes such as chloroform (CHCl_3), bromodichloromethane (CHBrCl_2), dibromochloromethane (CHBr_2Cl) and bromoform (CHBr_3) during the chlorination process has become evident (1, 2).

The United States Environmental Protection Agency conducted a comprehensive survey of 80 water supplies in the U.S. for halogenated organics (3). It was found that the four trihalomethanes—

CHCl_3 , CHBrCl_2 , CHBr_2Cl , CHBr_3 —are widespread in chlorinated drinking water in the U.S. and result from chlorination, since these substances were not found in raw waters. The trihalomethanes were not a result of impurities in the chlorine being used but were a result of the reaction of chlorine with precursor. Morris (4) made the point that the number of trihalomethane precursors is extensive and includes ethanol, acetaldehyde and the methyl ketones. Rook (1) and Morris (4) indicated that the bromide ion present in a natural water is rapidly oxidized by aqueous chlorine to hypobromous acid (HOBr), and that this acid reacts with precursor to account for the formation of the brominated species.

Since these trihalomethanes may be a health hazard, chlorine dioxide (ClO_2) is one of the alternative disinfectants being considered because it prevents the formation of trihalomethanes (5, 6). ClO_2 is a more effective oxidizing agent than chlorine and greatly improves the taste of water by avoiding the unpleasant taste associated with chlorination of

* Department of Pharmacology, UMDNJ-New Jersey Medical School, Newark, New Jersey 07103.

† Department of Pharmacology, College of Medicine, Ohio State University, Columbus, Ohio 43210.

†† Health Effects Research Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio 45268.

water (7). The unpleasant taste of chlorine in water is caused partially by the build-up of chlorinated phenols.

Miltner (6) has reported that the primary products resulting from ClO_2 disinfection of surface waters include chlorites (ClO_2^-) and chlorates (ClO_3^-), which appear in concentrations of 50% and 30% of ClO_2 demand, respectively.



Toxicity studies demonstrated that rats drinking ClO_2 , ClO_2^- and ClO_3^- (Cl compounds) daily for 9 months exhibited depressed red blood cell counts, hemoglobin concentration and packed cell volumes. Also, these Cl compounds in drinking water for three months inhibited the incorporation of ^3H -thymidine into nuclei of rat testes (8).

The studies described in this report were conducted to provide information on metabolism and pharmacokinetics of chlorine, chlorine dioxide, chlorite and chlorate in the rat. Also, the effect of ClO_2 in drinking water on the formation of CHCl_3 in rat blood and various organs was studied.

Methods

Synthesis of ^{36}Cl Compounds

H^{36}Cl was obtained from New England Nuclear. This material, with a radionuclidic purity of $> 99\%$ was used as the source of ^{36}Cl . The potassium chlorate ($\text{K}^{36}\text{ClO}_3$) was synthesized according to the method of Abdel-Rahman et al. (9). The generation of $^{36}\text{ClO}_2$ from $\text{K}^{36}\text{ClO}_3$ was accomplished by the reaction (1). $^{36}\text{Cl}_2$ was formed by the reaction (2).



The purity of the synthesized $\text{K}^{36}\text{ClO}_3$, which is an essential intermediate step to the production of ClO_2 , was 95%, as determined by fast neutron activation analysis.

Kinetics of Chlorine Dioxide, Chlorite, Chlorate and Chlorine in the Rat

Four groups of four male Sprague-Dawley rats (230 ± 20 g) each were used in these experiments. ^{36}Cl compounds were administered by gavage as follows: 3 ml of 100 mg/l. $^{36}\text{ClO}_2$ (0.7 μCi) was administered to each rat, producing a dose of 1.5 mg/kg body weight; 3 ml of 10 mg/l $^{36}\text{ClO}_2^-$ (0.17 μCi)

was administered to each rat, producing a dose of 0.15 mg/kg body weight; 3 ml of 5 mg/l. $^{36}\text{ClO}_3^-$ (0.85 μCi) was administered to each rat, producing a dose of 0.065 mg/kg body weight; 3 ml of 250 mg/l. HO^{36}Cl (0.45 μCi) was administered to each rat, producing a dose of 3.26 mg/kg body weight.

Heparinized blood samples were collected at 5, 10, 20, 30 and 60 min, and 2, 4, 8, 24 and 48 hr by orbital sinus puncture. The blood was centrifuged at 1000g for 15 min to separate the red blood cells from the plasma. Then the radioactivity was counted in a liquid scintillation counter (Beckman LS 7500), and the rate constant and $T_{1/2}$ of absorption and elimination from plasma were calculated for each compound.

Distribution of Cl Compounds in the Rat

After 72 hr following the administration of ^{36}Cl compounds as described previously, rats were killed by decapitation, and blood was collected in heparinized tubes. Tissue specimens of stomach, testes, lung, kidney, duodenum, ileum, spleen, liver, bone marrow, carcass and skin were prepared for the determination of ^{36}Cl content by liquid scintillation spectrometry as follows: Whole blood and packed cells were prepared according to the method of Mahin et al. (10). Samples of wet tissue weighing up to 200 mg and samples of bone marrow up to 50 mg were placed in individual glass LSC vials with Teflon or polyethylene-lined caps. Then 2 ml Protosol was added to each vial, which was capped tightly to prevent loss of solvent. Samples were solubilized by heating overnight at 50°C, then cooled at room temperature; afterwards 0.2 ml of 30% H_2O_2 was added for decolorization. The samples were heated at 50°C for 30 min, then cooled, and 15 ml of Aquasol-2 was added.

Sample quench effects were corrected by using the method of channel ratios, and efficiency was greater than 80%.

Metabolism and Excretion Studies

Four groups of four male Sprague-Dawley rats/group were used in these studies. Each group was administered orally the same concentrations of ^{36}Cl compounds described above. Animals were housed in modified Roth all-glass metabolism chambers for the collection of expired air and fecal and urine samples. The radioactivity of the total ^{36}Cl compounds was measured. The analysis of radioactive metabolites (Cl^- , ClO_2^- , ClO_3^-) was performed as described in a previous report, (9), and measured as a percentage of the initial dose.

Effect of Chlorine Dioxide, Chlorite and Chlorate on the Formation of Chloroform in Rat Blood and Organs

Male Sprague-Dawley rats (50–170 g) were used in this experiment. The animals were housed in a controlled environment (temperature, 22°C; humidity 50%) with a 12 hr light/dark cycle beginning at 6:00 A.M. Food was available *ad libitum*. Groups of rats were allowed to drink doubly distilled water containing 10 or 100 mg/l. ClO₂, ClO₂⁻ or ClO₃⁻ for 20 hr/day, 7 days/week for 12 months. ClO₂ was generated daily and water concentrations determined by the DPD (diethyl-*p*-phenylenediamine) method of Palin (11).

After one year, rats were killed by decapitation and blood collected in heparinized tubes. Tissue samples of liver, kidney, spleen, brain, and testes were homogenized in isotonic sucrose medium. Tissue homogenate or blood sample (3 ml) was extracted by 5 ml pentane. The pentane layer was then separated by centrifugation at 1000g for 5 min. A gas chromatograph equipped with an electron capture detector was used for the quantitation of chloroform in each sample, with bromodichloromethane being used as the internal standard.

Results

Kinetics of Cl Compounds in the Rat

The rate constant and $T_{1/2}$ for absorption and elimination from plasma of Cl compounds are shown in Table 1. The rate constant for absorption was highest for ³⁶ClO₂ (3.77 ± 0.24/hr), followed by ³⁶ClO₃⁻, ³⁶ClO₂⁻, and lowest for HO³⁶Cl (0.157 ± 0.001/hr). The $T_{1/2}$ for absorption, which is inversely proportional to the rate constant, was longest for HO³⁶Cl (4.42 ± 1.31 hr), followed by ³⁶ClO₂⁻, ³⁶ClO₃⁻, and was shortest for ³⁶ClO₂ (0.18 ± 0.01 hr). For elimination from plasma, the rate constant was highest for ³⁶ClO₂⁻ (0.02 ± 0.002/hr), followed by ³⁶ClO₃⁻, ³⁶ClO₂ and HO³⁶Cl (0.009 ± 0.001/hr). The $T_{1/2}$ elimination from plasma was longest for HO³⁶Cl (77.0 ± 8.8 hr), followed by ³⁶ClO₂, ³⁶ClO₃⁻, and ³⁶ClO₂⁻ (35.2 ± 3.0 hr).

Distribution of Cl Compounds in the Rat

The distribution of ³⁶Cl compounds after 72 hr from the time of oral administration is shown in Table 2. Values are expressed as a percentage of

Table 1. Kinetic constants of chlorine dioxide, chlorite, chlorate and chlorine in the rat.^a

Treatment	Absorption, hr ⁻¹		Elimination, hr ⁻¹	
	Rate constant	$T_{1/2}$	Rate constant	$T_{1/2}$
³⁶ ClO ₂ (100 mg/l.)	3.77 ± 0.24	0.18 ± 0.01	0.0158 ± 0.0008	43.9 ± 2.3
³⁶ ClO ₂ ⁻ (10 mg/l.)	0.198 ± 0.06	3.50 ± 1.06	0.0197 ± 0.0017	35.2 ± 3.0
³⁶ ClO ₃ ⁻ (5 mg/l.)	0.399 ± 0.151	1.74 ± 0.66	0.019 ± 0.003	36.7 ± 5.8
HO ³⁶ Cl (250 mg/l.)	0.157 ± 0.001	4.42 ± 1.31	0.009 ± 0.001	77.0 ± 8.8

^aValues were calculated from the time course of the elimination from rat plasma; *n* = 4 rats per treatment.

Table 2. Chlorine dioxide, chlorite, chlorate and chlorine distribution in the rat.

Tissue	Percentage of initial dose ^a			
	³⁶ ClO ₂ (100 mg/l.)	³⁶ ClO ₂ ⁻ (10 mg/l.)	³⁶ ClO ₃ ⁻ (5 mg/l.)	HO ³⁶ Cl (250 mg/l.)
Plasma	0.72 ± 0.02	0.55 ± 0.038	0.68 ± 0.09	0.77 ± 0.04
Packed cells	— ^b	0.63 ± 0.11	0.23 ± 0.02	—
Whole blood	—	0.64 ± 0.01	0.57 ± 0.05	—
Kidney	0.81 ± 0.15	0.30 ± 0.06	0.42 ± 0.07	0.39 ± 0.03
Lung	0.74 ± 0.15	0.37 ± 0.04	0.45 ± 0.07	0.34 ± 0.02
Stomach	0.70 ± 0.15	0.43 ± 0.07	0.46 ± 0.05	0.20 ± 0.03
Deodenum	0.29 ± 0.07	0.31 ± 0.01	0.34 ± 0.04	0.28 ± 0.04
Ileum	0.48 ± 0.09	0.17 ± 0.03	0.21 ± 0.02	0.14 ± 0.02
Liver	0.38 ± 0.09	0.06 ± 0.03	0.20 ± 0.03	0.20 ± 0.01
Spleen	0.25 ± 0.04	0.22 ± 0.02	0.29 ± 0.04	0.23 ± 0.02
Bone marrow	0.16 ± 0.03	0.09 ± 0.03	0.15 ± 0.03	0.40 ± 0.01
Testes	—	0.39 ± 0.04	0.45 ± 0.07	0.37 ± 0.02
Skin	—	0.38 ± 0.06	0.42 ± 0.10	0.32 ± 0.03
Carcass	—	0.25 ± 0.04	0.21 ± 0.02	0.18 ± 0.003

^aValues represent mean ± SE as percentage of the initial dose from four rats per treatment after 72 hr.

^bNot determined.

Table 3. Metabolism of chlorine dioxide, chlorite and chlorate in rat urine.

Treatment	Percentage of initial dose ^a		
	Cl ⁻	ClO ₂ ⁻	ClO ₃ ⁻
³⁶ ClO ₂ (100 mg/l.)	26.93 ± 1.6	3.46 ± 1.0	0.73 ± 0.73
³⁶ ClO ₂ ⁻ (100 mg/l.)	31.55 ± 4.5	6.0 ± 0.6	— ^b
³⁶ ClO ₃ ⁻ (5 mg/l.)	20.50 ± 0.7	3.95 ± 2.7	8.2 ± 2.9

^aValues represent the mean ± SE as percentage of the initial dose from four rats per treatment, throughout the 72 hr studied.

^bNone detected.

the initial dose. The distribution of ³⁶ClO₂ was highest in the kidney, followed by lung, plasma, stomach, ileum, liver, duodenum, spleen and bone marrow, while the distribution of HO³⁶Cl was highest in plasma, followed by bone marrow, kidney, testes, lung, skin, duodenum, spleen, stomach, liver, carcass and ileum. The distribution of ³⁶ClO₂⁻ was highest in plasma, followed by stomach, testes, skin, lung, duodenum, kidney, carcass, spleen, ileum, bone marrow and liver. In the case of ³⁶ClO₃⁻, plasma was highest, followed by stomach, lung, testes, kidney, skin, duodenum, spleen, ileum, carcass, liver and bone marrow.

In whole blood, when plasma was separated from packed cells, radioactivity was divided more or less

evenly between the plasma and packed cells (0.55 ± 0.038 and 0.63 ± 0.11% of initial dose, respectively) for ³⁶ClO₂⁻, whereas for ³⁶ClO₃⁻, radioactivity preferentially accumulated in the plasma (0.68 ± 0.09% of initial dose vs. 0.23 ± 0.02% in the packed cells).

Metabolism of Cl Compounds

The administered ³⁶Cl compounds are found as various metabolites in rat urine, as shown in Table 3. After 72 hr from the time of administration, ³⁶ClO₂ and ³⁶ClO₃⁻ are found in urine as Cl⁻, ClO₂⁻ and ClO₃⁻ whereas ³⁶ClO₂⁻ is found as Cl⁻ and ClO₂⁻. In all three cases, Cl⁻ is the major metabolite (approximately 87, 84 and 64% of the total radioactivity found in urine for ³⁶ClO₂, ³⁶ClO₂⁻ and ³⁶ClO₃⁻, respectively).

Excretion of Cl Compounds

The total of the ³⁶Cl compounds recovered in the urine and feces 72 hr after administration was determined for each compound and calculated as the percentage of the initial dose. In the case of ³⁶ClO₂, 75% of the recovered dose was found in the urine, while 25% was found in the feces. For ³⁶ClO₂⁻, 87 and 13% was found in urine and feces, respectively; and for HO³⁶Cl, 76 and 24% was found in urine and feces, respectively (Table 4).

Table 4. Excretion of chlorine dioxide, chlorite, chlorate and chlorine in the rat.

Treatment	Percentage of initial dose ^a	
	Urine	Feces
³⁶ ClO ₂ (100 mg/l.)	30.81 ± 0.8	10.10 ± 1.7
³⁶ ClO ₂ ⁻ (10 mg/l.)	34.51 ± 1.4	4.75 ± 1.0
³⁶ ClO ₃ ⁻ (5 mg/l.)	40.14 ± 2.14	3.14 ± 1.0
HO ³⁶ Cl (250 mg/l.)	21.52 ± 2.51	7.09 ± 0.24

^aValues represent the mean ± SE as percentage of the initial dose from four rats per treatment, throughout the 72 hr studied. ³⁶Cl was not detected in expired air.

Table 5. Effect of chlorine dioxide, chlorite and chlorate on the formation of chloroform in rat blood and organs.

Treatment ^a	Concentration, mg/l.	n	Organ	Effects
ClO ₂	10	4	Blood	No effect
	100	7	Blood	Increased
	10	4	Liver, kidney, spleen, brain	No effect
	10	4	Testes	Increased
	100	7	Liver, testes, brain	Increased
ClO ₂ ⁻	100	7	Blood	No effect
	100	7	Liver, brain	Increased
ClO ₃ ⁻	100	9	Blood	No effect
	100	9	Liver	Increased

^aRats were treated with chlorine dioxide and its metabolites daily for one year.

^{36}Cl compounds were not found in the expired air of any treatment group throughout the 72 hr time period.

Effect of Cl Compounds on the Formation of Chloroform in Rat Blood and Organs

Table 5 shows the effects of chronic treatment with chlorine dioxide and its metabolites for one year on the formation of chloroform in rat blood, liver, kidney, spleen, brain and testes. A concentration of 100 mg/l. ClO_2 daily in the drinking water caused an increase in chloroform levels in blood, liver, testes and brain. A concentration of 10 mg/l. ClO_2 increased chloroform levels in rat testes, while levels in blood, liver, kidney, spleen and brain remained unchanged. Treatment of rats with 100 mg/l. ClO_2^- for one year elevated chloroform levels in the liver and brain, while the levels in blood were not affected. Treatment with 100 mg/l. ClO_3^- increased liver but not blood chloroform concentrations after one year.

Discussion

From the kinetics study, it is apparent that ClO_2 and its metabolites (ClO_2^- and ClO_3^-) are more rapidly absorbed into the bloodstream than HOCl. The effect is very pronounced when comparing ClO_2 itself with HOCl. Likewise, HOCl is more slowly eliminated from rat plasma than ClO_2 or its metabolites.

The distribution study revealed that although plasma and whole blood contain the highest activity, the bone marrow contained far less in $^{36}\text{ClO}_2$, $^{36}\text{ClO}_2^-$ and $^{36}\text{ClO}_3^-$ treatment, while HOCl had a higher activity in bone marrow. The high activity in testes, even after 72 hr, indicates a possible pharmacologic action at this site. Effects of these compounds on the reproductive system are currently in progress in our laboratory.

It is important to indicate that after 72 hr, high amounts of radioactivity were still found in the stomach, duodenum and ileum.

The Cl compounds studied are eliminated mainly by kidney and intestinal routes, since no radioactivity was found in expired air. For HO^{36}Cl and $^{36}\text{ClO}_2$ treatment, about 75% of the recovered dose was found in the urine; in the case of $^{36}\text{ClO}_2^-$ and $^{36}\text{ClO}_3^-$, about 90% of the recovered dose was in the urine.

REFERENCES

1. Rook, J. J. Formation of haloforms during chlorination of natural waters. *J. Water Treat. Exam.* 23: 234-243 (1974).
2. Rook, J. J. Haloforms in drinking water. *J. Am. Water Works Assoc.* 68: 168-172 (1976).
3. Symons, J. M., Bellar, T. A., Carswell, J. K., DeMarco, J., Kropp, K. L., Robeck, G. G., Seeger, D. R., Slocum, C. J., Smith, B. L., and Stevens, A. A. National organics reconnaissance survey for halogenated organics in drinking water. *J. Am. Water Works Assoc.* 67: 634 (1975).
4. Morris, J. C. Formation of halogenated organics by chlorination of water supplies. U.S. EPA, Cincinnati, 1975, EPA 600/1-74-002.
5. Stevens, A., Seeger, D., and Slocum, C. J. Products of chlorine dioxide treatment of organic material in water. Paper presented at Workshop on Ozone Chlorine Dioxide Oxidation Products of Organic Materials, Cincinnati, Ohio, Water Supply Research Division, U.S. EPA, November 17-19, 1976.
6. Miltner, R. J. The effect of chlorine dioxide on trihalo-methanes in drinking water. Master Thesis, University of Cincinnati, Cincinnati, Ohio, 1976.
7. Enger, M. Treatment of water with chlorine dioxide to improve the taste. *Gass Wasser Fach* 14: 330-336 (1960).
8. Abdel-Rahman, M. S., Couri, D., and Bull, J. R. Toxicity of chlorine dioxide in drinking water. *J. Environ. Pathol. Toxicol.*, in press.
9. Abdel-Rahman, M. S., Couri, D., and Jones, J. Chlorine dioxide metabolism in rat. *J. Environ. Pathol. Toxicol.* 3 (1, 2): 421-430 (1979).
10. Mahin, D., and Lofberg, R. A simplified method of sample preparation for determination of ^3H , ^{14}C or ^{35}S in blood or tissue. *J. Anal. Biochem.* 16: 500-509 (1966).
11. Palin, A. T. Methods for the determination, in water, of free and combined available chlorine, chlorine dioxide and chlorite, bromine, iodine and ozone, using diethyl-*p*-phenylene diamine (DPD). *J. Inst. Water Engrs.* 21: 537-547 (1987).